

Data were assembled with care but treat differences below LSD cautiously. Performance variation is not AHDB's responsibility. AHDB Harvest Results 2017

AHDB Recommended Lists

Winter Triticale DL 1-year results 2017 - DL varieties

29-Nov Grain yield (t/ha) 2017 - treated - as % mean of controls				
Name	Code	2017 mean	Plumber Dorset	Clinge Sufflk
Control mean		9.92	9.28	10.55
LSD 5%		1.3	1.0	6.1
CV%		5.8	5.7	3.6
Agostino (C)	WT93	95	93	97
KWS Fido (C)	WT99	105	108	104
Tradiro	WT100	111	118	106
Adverdo	WT104	105	108	102
Kasyno	WT107	115	121	110
Dometica	WT108	110	113	107
Tribeca	WT9056	98	96	100
Cyrkon	WT9096	107	116	100
Kereon	WT9107	101	115	89
Securo	WT9117	101	109	94
LD17	WT9118	103	104	101

(C) = yield control

Not all varieties on the AHDB Recommended List are presented, since the focus is to test new varieties. Yields for well-established varieties on the AHDB Recommended Lists are available at cereals.ahdb.org.uk/varieties

The AHDB Recommended Lists are managed by a project consortium of AHDB Cereals & Oilseeds, BSPB, nabim and MAGB. AHDB Cereals & Oilseeds is a division of the Agriculture and Horticulture Development Board.

While the Agriculture and Horticulture Development Board seeks to ensure that the information contained within this document is accurate at the time of printing, no warranty is given in respect thereof and, to the maximum extent permitted by law, the Agriculture and Horticulture Development Board accepts no liability for loss, damage or injury howsoever caused (including that caused by negligence) or suffered directly or indirectly in relation to information and opinions contained in or omitted from this document. © Agriculture and Horticulture Development Board 2017. All rights reserved.

Harvest results from an individual trial sites should not be used to make conclusions on individual variety performance.

Recommended lists are based on information taken from trials across the UK over several seasons. As more information becomes available from the 2017 harvest, this will contribute to the over trials averages, which comprise 4 years for oilseeds and 5 years for cereals. These values provide better information for comparison between varieties.

When more results are available, our validation procedures will examine cross site differences further.